arXiv:1404.5294v1 [math.RA] 22 Apr 2014

ON THE QUIVER WITH RELATIONS OF A QUASITILTED ALGEBRA AND APPLICATIONS

NATALIA BORDINO, ELSA FERNÁNDEZ, AND SONIA TREPODE

Abstract. In this paper we discuss, in terms of quivers with relations, sufficient and necessary conditions for an algebra to be a quasitilted algebra. We start with an algebra with global dimension two and we give a sufficient condition for it to be a quasitilted algebra. We show that this condition is not necessary. In the case of a strongly simply connected schurian algebra, we discuss necessary conditions, and, combining both type of conditions, we are able to analyze if the given algebra is quasitilted. As an application we obtain the quiver with relations of all the tilted and cluster tilted algebras of Dynkin type E_p.

1. Introduction

An interesting problem in the theory of representation of algebras is to know if an algebra given by a quiver with relations is a tilted algebra. This problem was studied by several authors and has been solved in particular cases, see [1], [10], [11], [12], [19], [20], [21], [16]. In this work, we consider a bigger class of algebras, that is, the quasitilted algebras introduced by Happel, Reiten and Smalo, see [17].

We introduced a notion of bounded consecutive relations inspired by the one given by Assem and Redondo in [3]. Our first result states that, an algebra with global dimension two having non consecutive bounded relations, is a quasitilted algebra. Therefore, we show that the converse of this result does not hold. On the other hand, if a quasitilted algebra has bounded consecutive relations, we give a necessary condition over this kind of relations.

Finally, we obtain a sufficient condition for a strongly simply connected schurian algebra to be a quasitilted algebra. We arrive at this result combining the criterion for global dimension two, developed in [6] with the previous results.

As applications of our main result, and using the results given in [13], we are able to give all the quivers with relations of the tilted and cluster tilted

2000 *Mathematics Subject Classification.* Primary: 16G20, Secondary: 16G70,

We thank M. I. Platzeck for useful discussions about Theorem 3.3. This work is part of the Ph.D. Thesis of Natalia Bordino, under the supervision of her advisers, Sonia Trepode and Elsa Fernández. The third author is a researcher of CONICET, Argentina.

algebras of Dynkin type. In this work we consider tilted an cluster tilted algebras of type E_p.

2. Preliminaries

In this paper, by an algebra, we always mean a basic and connected finite dimensional algebra over an algebraically closed field k. Given a quiver Q, we denote by Q_0 its set of vertices and by Q_1 its set of arrows. A *relation* in Q from a vertex x to a vertex y is a linear combination $\rho = \sum_{i=1}^{m} \lambda_i w_i$ where, for each i, $\lambda_i \in k$ is non-zero and w_i is a path of length at least two from x to y.

We denote by kQ the path algebra of Q and by $kQ(x, y)$ the k-vector space generated by all paths in Q from x to y. For an algebra A, we denote by Q_A its ordinary quiver. For every algebra A, there exists an ideal I in kQ_A, generated by a set of relations, such that $A \simeq kQ_A/I$. The pair (Q_A, I) is called a *presentation* of A and A is said to be given by the *bound quiver* (Q_A, I).

An algebra A is called *triangular* if Q_A has no oriented cycles, and it is called *schurian* if, for all $x, y \in A_0$, we have $\dim_k A(x, y) \leq 1$. A triangular algebra A is called *simply connected* if, for any presentation (Q_A, I) of A, the group $\pi_1(Q_A, I)$ is trivial, see [5]. It is called *strongly simply connected* if every full convex subcategory of A is simply connected, [23].

In this work, we always deal with schurian triangular algebras. For a vertex x in the quiver Q_A, we denote by e_x the corresponding primitive idempotent, S_x the corresponding simple A-module, and by P_x and I_x the corresponding indecomposable projective and injective A-module, respectively.

Let A be an algebra. A module T_A is called a *tilting module* [16] if $pdT_A \leq 1$; $\mathrm{Ext}^1_A(T; T) = 0$ and the number of isomorphism classes of indecomposable summands of T equals the rank of the Grothendieck group $K_0(A)$. An algebra A is called *tilted of type* Q if it is the endomorphism algebra of a tilting kQ-module.

An algebra A is called *quasitilted* if gl.dim. $A \leq 2$ and, for each indecomposable module M_A, we have $pd\, M \leq 1$ or $id\, M \leq 1$ (see [17]). It follows from [17], that tilted algebras are a subclass of quasitilted algebras, and if a representation finite algebra is quasitilted, then it is tilted.

3. On the relations of a quasitilted algebra

The main objective of this section is the interaction between the relations of a quasitilted algebra. More precisely, for this class of algebras we are going to study how the relations can interact.

We begin by introducing the concept of bound consecutive relations inspired by the ones introduced in [3]. Observe that the pair of relations defined in [3] is a subclass of the bound consecutive relations defined in the following.

Definition 3.1. The relations ρ_1 and ρ_2 are called *bound consecutive relations* if there is a walk ϖ between $t(\rho_1)$ and $s(\rho_2)$ such that ϖ does not contain zero subpaths.

For example, if $A = kQ/I$ is the algebra given by the quiver

Q



bound by $\alpha_5\alpha_1\alpha_3 = 0$ and $\alpha_5\alpha_1\alpha_6 = 0$. These relations are bound consecutive for the walk $\varpi = \alpha_4\alpha_2\alpha_5$:



The main theorem of this section, states that any algebra of global dimension two that does not contain bound consecutive relations is quasitilted.

Theorem 3.2. *Let A be an algebra with* gl.dim. $A \leq 2$. *If there are no bound consecutive relations in A, then A is quasitilted.*

Proof. Suppose A is not a quasitilted algebra. Then, there is an indecomposable A-module M such that $pd\, M = id\, M = 2$.

Let

$$0 \longrightarrow P_2 \xrightarrow{f_2} P_1 \xrightarrow{f_1} P_0 \xrightarrow{f_0} M \longrightarrow 0$$

and

$$0 \longrightarrow M \longrightarrow I_0 \longrightarrow I_1 \longrightarrow I_2 \longrightarrow 0$$

be the minimal projective and injective resolutions of M, respectively.

Any indecomposable sum $P_2(S_b)$ of P_2 gives rise to a diagram $P_2(S_b) \longrightarrow P_1 \longrightarrow P_0$ with zero composition. Then there is a sum $P_0(S_j)$ of P_0 and a relation $P_2(S_b) \longrightarrow P_1 \longrightarrow P_0(S_j)$. Consequently, there is a relation $\rho_2 : j \rightsquigarrow b$ starting at the vertex j, where $S_j \in \operatorname{Top} M$.

Dually, using the minimal injective resolution of M, there is $S_i \in \operatorname{Soc} M$ and a relation ρ_1 ending at the vertex i,

Let $S_k \in \operatorname{Top} M$ such that $S_i \in f_0(P_0(S_k))$. Then there exists a nonzero morphism $h : P_0(S_i) \longrightarrow f_0(P_0(S_k))$ and an epimorphism $g : P_0(S_k) \longrightarrow f_0(P_0(S_k))$, i.e. the situation is as follows:



Then there is a nonzero morphism $f : P_0(S_i) \longrightarrow P_0(S_k)$ such that $gf = h$. Therefore, there is a nonzero path $\gamma : k \rightsquigarrow i$.

Since M is an indecomposable module, then $\operatorname{sop}M$ is connected. We consider $Q'_0 = (\operatorname{sop} M)_0$ and $Q'_1 = \{\alpha \in Q_1 \text{ such that } M(\alpha) \neq 0\}$. Consequently, $Q' = (Q'_0, Q'_1)$ is a connected quiver and the vertices associated with the simple A-modules of $\operatorname{Top} M$ are sources of Q'. Given two sources j and k, there is a walk of nonzero paths that unites them, ([4], pp. 45). Therefore, ρ_1 y ρ_2 are bound consecutive relations □

We want to remark that the algebra A, given in the previous example, is not quasitilted.

We illustrated with the following example that the reciprocal of Theorem 3.2 is generally not true. The algebra B given by the following quiver with relations:



is a tilted algebra and the given relations are bound consecutive relations, where $\varpi : \ 1 \longleftarrow 2 \longleftarrow 4$ is the path.

Next, we give necessary conditions for an algebra of global dimension two to be quasitilted. To this end, we introduce the following notations.

Let $A = kQ/I$ be an algebra and $\rho = \sum_{i=1}^{m} \lambda_i \alpha_{i_1} \alpha_{i_2} \dots \alpha_{i_{r_i}}$ be a relation of A, with $\alpha_{i_j} \in Q_1$. We will note with $\mathbf{Q}_\rho$ the subquiver of Q induced by the arrows of ρ, i.e. $(\mathbf{Q}_\rho)_1 = \{\alpha_{i_j} : 1 \leq i \leq m, 1 \leq j \leq r_i\}$ and $(\mathbf{Q}_\rho)_0 = \{s(\alpha_{i_j}) : \alpha_{i_j} \in (\mathbf{Q}_\rho)_1\} \cup \{t(\alpha_{i_j}) : \alpha_{i_j} \in (\mathbf{Q}_\rho)_1\}$.

If $A = kQ/I$ is a schurian strongly simply connected algebra and ρ_1, ρ_2 are bound consecutive relations for the walk

$$\varpi = \gamma_1 \gamma_2 \cdots \gamma_r : t(\rho_1) = i \rightsquigarrow s(\rho_2) = j$$

with nonzero maximal paths γ_h and $\text{long}(\gamma_h) \neq 0$, for $h = 2, \ldots, r-1.$, then we note with $\overline{\text{sop}}\, \varpi$ the support of the union of all the parallel paths of γ_h (which are all equal and nonzero in A).

The following result has a strong relation with Lemma 2.1, given by I. Assem and M. J. Redondo in [3]. They proved that a schurian tilted algebra A does not contain a certain class of bound consecutive relations. In the case of schurian strongly simply connected quasitilted algebras, we show that these algebras do not have the bounded consecutive relations defined in this work.

Theorem 3.3. *Let $A = kQ/I$ be a schurian strongly simply connected algebra with* gl.dim. $A = 2$. *If A is quasitilted, then there are no bound consecutive relations ρ_1 y ρ_2 with walk ϖ such that $\overline{\text{sop}}\, \varpi \cap \mathbf{Q}_{\rho_1} = \{t(\rho_1) = i\}$ and $\overline{\text{sop}}\, \varpi \cap \mathbf{Q}_{\rho_2} = \{s(\rho_2) = j\}$.*

Proof. Since A is schurian strongly simply connected, we can define a representation M whose support is $\overline{\text{sop}}\, \varpi$ and such that $M(x) = k$, for all $x \in (\text{sop}\, M)_0$ and $M(\varepsilon) = Id_k$, for all $\varepsilon \in (\text{sop}\, M)_1$. We will show that the projective dimension of M is 2.

Consider

$$P_1(M) \xrightarrow{f_1} P_0(M) \xrightarrow{f_0} M \longrightarrow 0$$

The module M is not projective because $P_j \in \text{add}P_0(M)$ and $\text{sop}P_j \nsubseteq \text{sop}M$.

If $\rho_2 = \beta\delta_2$, where $\beta \in Q_1$ and δ_2 is a nonzero path in A, then $f_0(\beta) = 0$ because $\overline{\text{sop}}\, \varpi \cap \mathbf{Q}_{\rho_2} = \{s(\rho_2) = j\}$. Then, $P_{t(\beta)}$ is a direct summand of $P_1(M)$ because the composition $P_{t(\beta)} \xrightarrow{\beta\cdot} P_j \xrightarrow{f_0} M$ is zero. Therefore, $Im\, \beta\cdot \subseteq \text{Ker}\, f_0$.

As $\beta \notin \text{rad Ker}\, f_0$, we have that $Im\, \beta\cdot \nsubseteq \text{rad Ker}\, f_0$. This proves that $P_{t(\beta)} \xrightarrow{\beta\cdot} P_j$ is a sum of the projective presentation of M.

Finally, as $0 = \rho_2 = \beta\delta_2 = \beta \cdot (\delta_2)$, it follows that $0 \neq \delta_2 \in \text{Ker}\ \beta\cdot$. Consequently, $P_2(M) \neq 0$.

If $\rho_2 = \beta\delta_2 + \beta'\delta_2'$, where $\beta, \beta' \in Q_1$ and δ_2, δ_2' are nonzero paths in A. Then, analogously to the previous case proof that $P_{t(\beta)} \oplus P_{t(\beta')} \xrightarrow{(\beta\cdot, \beta'\cdot)} P_j$ is a sum of projective presentation of M. Then, $0 = \rho_2 = \beta\delta_2 + \beta'\delta_2' = (\beta\cdot, \beta'\cdot)\begin{pmatrix} \delta_2 \\ \delta_2' \end{pmatrix}$. Therefore, $P_2(M) \neq 0$.

By duality, considering the minimal injective co-resolution of M, it follows that $id\, M = 2$. □

Example 3.4. Let $A = kQ/I$ be the algebra given by:



The two relations of this algebra are minimal bound consecutive relations with a walk $\varpi : \ 4 \longleftarrow 5$:

As the A-module M, whose representation is:



has projective dimension and injective dimension equal to 2. Then, by the previous theorem, A is not a quasitilted algebra.

The next question is to study what happens when a quasitilted algebra has a bound consecutive relation by a walk ϖ such that $\overline{\text{sop}}\ \varpi$ intersects the quiver of one of the relations in at least one arrow. In the next theorem we answer the question posed above, to schurian strongly simply connected algebras.

Theorem 3.5. *Let $A = kQ/I$ be a schurian strongly simply connected algebra and quasitilted such that there are ρ_1 and ρ_2 bound consecutive relations by the walk ϖ. Then, either $\overline{\text{sop}}\ \varpi \cap Q_{\rho_1}$ or $\overline{\text{sop}}\ \varpi \cap Q_{\rho_2}$ is a maximal subpath of ρ_1 or ρ_2, respectively.*

Proof. Suppose that for $l = 1, 2$, $\overline{\text{sop}}\ \varpi$ and Q_{ρ_l} do not intersect in a maximal subpath of ρ_l. Let M be the representation whose support is $\overline{\text{sop}}\ \varpi$ and such that $M(x) = k$, for all $x \in (\text{sop}\, M)_0$ and $M(\varepsilon) = Id_k$, for all $\varepsilon \in (\text{sop}\, M)_1$. If we prove that the projective dimension of M is 2, then by duality $di\, M = 2$.

As M is not a projective module, we can consider

$$0 \neq P_1(M) \xrightarrow{f_1} P_0(M) \xrightarrow{f_0} M \longrightarrow 0.$$

For $\rho_2 = \delta_2\beta$, where $\beta \in Q_1$ and δ_2 is a nonzero path in A, must be $f_0(\delta_2) = 0$. Writing $\delta_2 = \psi_2\varphi_2$, with ψ_2 minimal path in δ_2 such that $f_0(\psi_2) = 0$, results that the composition $P_{t(\psi_2)} \xrightarrow{\psi_2 \cdot} P_j \xrightarrow{f_0} M$ is zero. Therefore, $Im\, \psi_2 \cdot \subseteq \text{Ker}\ f_0$.

Suppose that $\psi_2 \in \text{rad Ker} f_0$. Then there exist non-trivial paths λ_l such that $\psi_2 = \sum \mu_l\lambda_l$, with $\mu_l \in \text{Ker}\ f_0$. As A schurian strongly simply connected,

$\psi_2 = a\lambda_1\mu_1$, with a scalar. But $f_0(\mu_1) = 0$ and $l(\mu_1) < l(\psi_2)$, which contradicts the minimality of ψ_2. Then, $Im\,\psi_2\cdot \nsubseteq \text{rad Ker}\, f_0$. Consequently, $P_{t(\psi_2)} \xrightarrow{\psi_2\cdot} P_j$ is a sum of projective presentation of M.

Since $0 = \rho_2 = \psi_2\varphi_2\beta = \psi_2 \cdot (\varphi_2\beta)$, Then $0 \neq \varphi_2\beta \in \text{Ker } \psi_2\cdot$. Therefore, $P_2(M) \neq 0$.

For $\rho_2 = \delta_2 + \delta_2'$, where δ_2, δ_2' are nonzero paths in A, it follows that $f_0(\delta_2) = f_0(\delta_2') = 0$. If we write $\delta_h = \psi_h\varphi_h$, with ψ_h minimal path in δ_h such that $f_0(\psi_h) = 0$, for $h = 1, 2$, is proved similarly to the previous case that $P_{t(\psi_2)} \oplus P_{t(\psi_2')} \xrightarrow{(\psi_2\cdot, \psi_2'\cdot)} P_j$ is a sum of projective presentation of M.

Finally, since $0 = \rho_2 = \delta_2 + \delta_2' = \psi_2\varphi_2 + \psi_2'\varphi_2' = (\psi_2\cdot, \psi_2'\cdot)\begin{pmatrix} \varphi_2 \\ \varphi_2' \end{pmatrix}$ we have that $P_2(M) \neq 0$. □

Example 3.6. Let $A = kQ/I$ be the tilted algebra given by the following quiver with relations:



Note that the two monomial relations having A are bound consecutive by the path $\varpi : 4 \longleftarrow 2 \longleftarrow 3$, that is:



However, $\overline{\text{sop}}\,\varpi$ intersects the relation $\rho : 3 \longrightarrow 2 \longrightarrow 1$ in the path $3 \longrightarrow 2$, which is maximal in ρ.

The condition of Theorem 3.5 is not a sufficient condition to ensure that one is quasitilted algebra. In fact, let A be the algebra given by the following quiver with relations:



and we will consider the following monomial relations: $\rho_1 = 7 \longrightarrow 4 \longrightarrow 5$ and $\rho_2 = 3 \longrightarrow 2 \longrightarrow 1$. Then, ρ_1 and ρ_2 are bound consecutive relations by the walk $\varpi : 5 \longleftarrow 4 \longrightarrow 2 \longleftarrow 3$:



It is clear that $\overline{\mathrm{sop}}\,\varpi$ intersects Q_{ρ_1} and Q_{ρ_2} in maximal subpaths of ρ_1 and ρ_2, respectively.

However, the A-module M given by the following representation



is indecomposable and $pd\,M = id\,M = 2$. Therefore, A is not quasitilted algebra.

4. On the quiver with relations of a quasitilted algebra

We study the quiver with relations of a quasitilted algebra. We start recalling the definition of a critical algebra given in [6].

Definition 4.1. Let B be an algebra. We say that B is *critical* if it satisfies the following properties:

i) B has a unique source i and a unique sink j,
ii) $pd\,S_i = id\,S_j = 3$ and if S is a different simple B-module we have that $pd\,S \leq 2$ and $id\,S \leq 2$,
iii) Let $0 \to \mathsf{Q}_3 \to \mathsf{Q}_2 \to \mathsf{Q}_1 \to \mathsf{Q}_0 \to S_i \to 0$ be the minimal projective resolution of the simple S_i and let $\mathsf{Q} = \oplus_{k=0}^{3}\mathsf{Q}_k$. Then all indecomposable projective B-modules are in $\mathrm{add}\mathsf{Q}$, and each indecomposable projective B-module is a direct summand of exactly one Q_k,
iv) Let $0 \to S_j \to I_0 \to I_1 \to I_2 \to I_3 \to 0$ be the minimal co-injective resolution of simple S_j and let $I = \oplus_{k=0}^{3} I_k$. Then all indecomposable injective B-modules are in $\mathrm{add}\,I$, and each indecomposable injective B-module is a direct summand of exactly one I_k.
v) B does not contain any proper full subcategory that verifies i), ii), iii) and iv).

We have the following corollary.

Corollary 4.2. *Let A be a strongly simply connected schurian algebra. If the following conditions hold:*

a) *A does not contain a proper full subcategory B such that B is a critical algebra.*
b) *A has no bounded consecutive relations.*

Then A is a quasitilted algebra.

Proof. It follows from a) and [6] that $\mathrm{gl.dim.}A \leq 2$. Appling Theorem 3.2 we get that A is quasitilted algebra. □

5. Applications

5.1. **Tilted algebras of Dynkin type.** In this section we provide a method to obtain a complete classification of tilted algebras of Dynkin type. In particular we give the classification of tilted algebras of type E_6.

The combinatorial description of how the iterated tilted algebra of Dynkin type can be obtained from a trivial extension of finite representation type was studied in [13]. One of the essential tools in this description is the notion of admissible cuts of trivial extensions (see [13] and [14]).

The next step is to select the iterated tilted algebras with global dimension at most two, using [6]. For this propose we use the Theorem 3.2, Theorem 3.3 and Theorem 3.5.

In this way, we could obtain a classification of the tilted algebras of Dynkin type. We now give the classification of Dynkin type E_6.

In the following, we will consider figures obtained from a quiver considering some arrows without orientation. Moreover, it is said that a quiver is associated with a figure if it is obtained giving an orientation to these arcs.

Theorem 5.1. *Let B be an algebra. Then B is tilted of Dynkin type E_6 if and only if*

i. *B is hereditary, or*
ii. *B or B^{op} is isomorphic to one of the following:*





The characterization of tilted algebras of type A_n in terms of their quiver with relations has been done by I. Assem in [1]. F. Huard, in [18], has characterized the tilted algebras kQ/I where the underlying graph of Q is D_n. An alternative proof of this result can be made using the method

described in this work. A complete classification of tilted algebras of type $E_p, p = 7, 8$, was done in [7].

5.2. **Cluster tilted algebras of finite representation type.** The relationship between tilted algebras and cluster-tilted algebras was given in [2]. For this, the authors introduced the concept of relation extension $R(B)$ of an algebra B with gl.dim.$B \leq 2$. Recall that the algebra $R(B) = B \propto \mathrm{Ext}^2_B(DB, B)$, where $DB = \mathrm{Hom}_k(B, k)$. In case the quiver of B has no oriented cycles, a construction of the quiver of the relation extension algebra is given in [2] . The main result in [2] characterizes the cluster tilted algebras as the relation-extension of tilted algebras.

On the other hand, A. Buan, R. Marsh and I.Reiten in [8] proved that any cluster tilted algebra of finite representation type is uniquely determined by its ordinary quiver (up to isomorphism).

Using Theorem 5.1 and the two previous results, we deduce the following Theorem.

Theorem 5.2. *Let C be an algebra. Then C is cluster tilted of Dynkin type E_6 if and only if*

i. *C is a hereditary algebra of Dynkin type E_6, or*
ii. *the quiver Q_C or $Q_{C^{op}}$ is isomorphic to one of the following:*



It is interesting to point out that, the classification of the cluster tilted algebras of the remaining Dynkin types, can be done in a similar way. Since the cluster tilted algebras of type A_n and D_n were studied in [9] and [22] respectively, by geometric methods, and in [15] was shown that it is not possible to obtain the E_p type using geometric methods, we are mainly interested in the cluster tilted algebras of type $E_p, p = 7, 8$. For classification of these two classes of algebras see [7].

References

[1] I. Assem. *Tilted algebras of type A_n*. Comm. Algebra 19, (1982), 2121-2131.

[2] I. Assem, T. Brüstle, R. Schiffler. *Cluster-tilted algebras as trivial extensions*. Bull. Lond. Math. Soc. 40 (2008), no. 1, 151-162.
[3] I. Assem, M. J. Redondo. *The first Hochschild cohomology group of a schurian cluster-tilted algebra*. Manuscripta Math. 128, (2009), no. 3, 373-388.
[4] M. Auslander, I. Reiten, S. Smalø. *Representation Theory of Artin Algebras*, Cambridge Studies in Advanced Mathematics, vol 36, Cambridge University Press (1995).
[5] K. Bongartz, P. Gabriel. *Covering Spaces in Representation-Theory*. Invent. Mathematicae 65, (1982), 331-378.
[6] N. Bordino, E. Fernández, S. Trepode. *A criterion for global dimension two for strongly simply connected schurian algebras*. arXiv:1110.6160 [math.RA].
[7] N. Bordino, E. Fernández, S. Trepode. *Quivers with relations of tilted algebras and cluster tilted algebras of type E_p*. In preparation.
[8] A. Buan, R. Marsh, I. Reiten. *Cluster-tilted algebras of finite representation type*. J. Algebra 306, (2006), no. 2, 412-431.
[9] P. Caldero, F. Chapoton, R. Schiffler. *Quivers with relations arising from clusters (A_n case)*. Trans. Amer. Math. Soc. 358, (2006), no. 3, 1347-1364.
[10] F. Coelho, J. A. de la Peña, S. Trepode. *Non-quasitilted til critical algebras*. XVI Coloquio Latinoamericano de Álgebra, Proceedings 2007.
[11] F. Coelho, J. A. de la Peña, S. Trepode. *Minimal non tilted algebras*. Coloquium Mathematicum 111, (2008), 71-81.
[12] F. Coelho, J. A. de la Peña, S. Trepode. *Tilt-critical algebras of tame type*. Comm. Algebra. Volume 36, (2008), no. 9, 3574-3584.
[13] E. Fernández. *Extensiones triviales y álgebras inclinadas iteradas*. Ph.D. Thesis, Universidad Nacional del Sur, Argentina (1999).
[14] E. Fernández, M. I. Platzeck. *Presentations of trivial extensions of finite dimensional algebras and a theorem of Sheila Brenner*. J. Algebra 249, (2002), 326-344.
[15] S. Fomin, M. Shapiro, D. Thurston. *Cluster algebras and triangulated surfaces. Part I: Cluster complexes*. Acta Math. 201, (2008), 83 - 146.
[16] D. Happel, C. Ringel. *Tilted algebras*. Trans. Amer. Math. Soc. 274, (1982), no. 2, 399-443.
[17] D. Happel, I. Reiten, S. O. Smalø. *Tilting in Abelian Categories and Quasitilted algebras*. Memoirs Amer. Math. Soc. 575, (1996).
[18] F. Huard. *Tilted algebras having underlying graph D_n*. Annales des Sciences Mathématiques du Québec (2001), 25 (1) (2001) 39-46.
[19] F. Huard. *Tilted gentle algebras*. Comm. Algebra 26, (1998), no. 1, 63-72.
[20] F. Huard, S. Liu. *Tilted special biserial algebras*. J. Algebra (1999), 679-700.
[21] F. Huard, S. Liu. *Tilted string algebras*. J. Pure Appl. Algebra 153, (2000), 151-164.
[22] R. Schiffler. *A geometric model for cluster categories of type D_n*. J. Algebraic Combin. 27, (2008), no. 1, 1-21.
[23] A. Skowroński. *Simply connected algebras and Hochschild cohomologies*. Representations of algebras (Ottawa, ON, 1992), 431–447, CMS Conf. Proc., 14, Amer. Math. Soc., Providence, RI, 1993.

Natalia Bordino, Departamento de Matemática, Facultad de Ciencias Exactas y Naturales, Funes 3350, Universidad Nacional de Mar del Plata, 7600 Mar del Plata, Argentina.

E-mail address: bordino@mdp.edu.ar

Elsa Fernández, Facultad de Ingeniería, Universidad Nacional de la Patagonia San Juan Bosco, 9120 Puerto Madryn, Argentina.

E-mail address: elsafer9@gmail.com

Sonia Trepode, Departamento de Matemática, Facultad de Ciencias Exactas y Naturales, Funes 3350, Universidad Nacional de Mar del Plata, 7600 Mar del Plata, Argentina.

E-mail address: strepode@mdp.edu.ar

arXiv:1404.5294v1 [math.RA] 22 Apr 2014

QUIVERS WITH RELATIONS OF TILTED ALGEBRAS AND CLUSTER TILTED ALGEBRAS OF TYPE E_P

NATALIA BORDINO, ELSA FERNÁNDEZ, AND SONIA TREPODE

Abstract. In this note we list the quiver with relations of all tilted and cluster tilted algebras of Dynkin type E_p.

The quiver with relations of all these algebras were obtained using the method developed in [1].

In this note, we will consider figures obtained from a quiver considering some arrows without orientation. Moreover, it is said that a quiver is associated with a figure if it is obtained by giving an orientation to these arcs.

1. Tilted algebras of type E_p

1.1. Quivers with relations of tilted algebras of type E_6.



2000 *Mathematics Subject Classification.* Primary: 16G20, Secondary: 16G70,

This work is part of the Ph.D. Thesis of Natalia Bordino, under the supervision of her advisers, Sonia Trepode and Elsa Fernández. The third author is a researcher of CONICET, Argentina.


5.4)
5.5)
6.1)
6.2)
6.3)
7.1)
7.2)
7.3)
8.1)
8.2)
8.3)
8.4)
8.5)
9.1)
9.3)
10.1)
10.2)
10.3)
10.4)
10.5)
10.6)
11.1)
11.2)
11.3)
11.4)
12.1)
12.2)
12.3)
12.4)
13.1)
13.2)
13.3)
13.4)
13.5)
14.1)
14.2)
14.3)
14.4)
14.5)
15.1)
15.2)
15.3)
15.4)
15.5)
15.6)
15.7)

16)

1.2. **Quivers with relations of tilted algebras of type E_7.**

1.1)

1.2)

1.3)



2.1)



2.2)



2.3)



3.1)



3.2)



3.3)

4.1)



4.2)



4.3)



4.4)



4.5)



4.6)

4.7)



5.1)



5.2)

5.3)

5.4)

5.5)



5.6)



5.7)



6.1)



6.2)

6.3)



6.4)

7.1)



7.2)



7.3)



7.4)



8.1)



8.2)

8.3)



8.4)



8.5)



9.1)




9.2)
9.3)
9.4)
9.5)
10.1)
10.2)
10.3)
10.4)
10.5)
11.1)
11.2)
11.3)
11.4)
11.5)
11.6)
12.1)
12.2)
12.3)
13.1)
13.2)
13.3)
14.1)
14.2)
14.3)
15.1)
15.2)
15.3)
16.1)
16.2)
16.3)
16.4)
16.5)
16.6)
16.7)
17.1)
17.2)
17.3)
17.4)

17.5) 

17.6) 

17.7) 

18.1) 

18.2) 

18.3) 

18.4)

18.5)

18.6)

18.7)

19.1) 

19.2) 

19.3) 

19.4) 

19.5) 

20.1) 

20.2) 

20.3) 

20.4) 

20.5) 

21.1) 

21.2) 

21.3) 

21.4) 

21.5) 

22.1) 

22.2) 

22.3) 

22.4) 

22.5) 

23.1) 

23.2) 

23.3) 

23.4) 

23.5)

24.1) 

24.2) 

24.3) 

24.4) 

24.5) 

25.1) 

25.2) 
25.3) 
25.4) 
25.5) 
25.6) 
26.1) 
26.2) 
26.3) 
26.4) 
26.5) 
26.6) 
27.1) 
27.2) 
27.3) 
27.4) 
27.5) 
27.6) 
27.7) 
27.8) 
28.1) 
28.2) 
28.3) 
28.4) 
28.5) 
28.6) 
28.7) 
28.8) 
29.1) 
29.2) 
29.3) 
29.4) 
29.5) 
29.6) 
29.7) 
29.8) 
29.9) 
29.10) 
29.11) 
30.1) 
30.2) 
30.3) 

30.4) 
30.5) 
30.6)
31.1) 
31.2) 
31.3) 
31.4) 
31.5) 
31.6) 
31.7) 
31.8) 
31.9) 
32.1) 
32.2) 
32.3) 
32.4) 
32.5) 
33.1) 
33.2) 
33.3) 
33.4) 
33.5)
33.6) 
33.7) 
33.8) 
33.9) 
34.1) 
34.2) 
34.3) 
34.4) 
34.5) 
34.6) 
34.7) 
34.8) 
34.9) 
34.10) 
34.11) 
35.1) 
35.2) 
35.3) 
35.4) 
35.5) 
35.6)
36.1) 
36.2) 
36.3) 
36.4) 
36.5) 


36.6)
36.7)
36.8)
37.1)
37.2)
37.3)
37.4)
37.5)
38.1)
38.2)
38.3)
38.4)
38.5)
39.1)
39.2)
39.3)
39.4)
39.5)
40.1)
40.2)
40.3)
41.1)
41.2)
41.3)
41.4)
41.5)
42.1)
42.2)
42.3)
42.4)
42.5)
43.1)
43.2)
43.3)

43.4) 

43.5)

44.1)

44.2)

44.3)

44.4) 

44.5) 

44.6) 

44.7) 

44.8) 

44.9) 

44.10) 

45.1) 

45.2)

45.3) 

45.4) 

45.5) 

45.6) 

45.7) 

46.1)

46.2) 

46.3) 

46.4) 

46.5) 

46.6) 

46.7 

47.1) 

47.2) 

47.3) 

47.4) 

47.5) 

47.6) 

47.7)

48.1) 

48.2) 

48.3) 

48.4) 

48.5) 

48.6) 

48.7) 


49.1)
49.2)
49.3)
49.4)
49.5)
49.6)
50.1)
50.2)
50.3)
50.4)
50.5)
50.6)
50.7)
50.8)
50.9)
50.10)
50.11)
50.12)
51.1)
51.2)
51.3)
51.4)
51.5)
51.6)
51.7)
51.8)
51.9)
51.10)
51.11)
52.1)
52.2)
52.3)
52.4)
52.5)
52.6)
52.7)
52.8)
53.1)
53.2)
53.3)
53.4)
53.5)
53.6)

53.7) 

53.8) 

54.1) 

54.2) 

54.3) 

54.4) 

54.5) 

54.6) 

54.7) 

54.8) 

55.1) 

55.2) 

55.3) 

55.4) 

55.5) 

56.1) 

56.2) 

56.3) 

56.4) 

56.5) 

56.6) 

56.7) 

56.8) 

56.9) 

56.10) 

56.11) 

56.12) 

57.1) 

57.2) 

57.3) 

57.4) 

57.5) 

57.6) 

58.1) 

58.2) 

58.3) 

58.4) 

58.5) 

58.6) 

58.7) 

58.8) 
58.9) 
59.1) 
59.2) 
59.3) 
59.4) 
59.5) 
59.6) 
59.7) 
59.8) 
59.9) 
60.1) 
60.2) 
60.3) 
60.4) 
60.5) 
60.6) 
60.7) 
61.1) 
61.2) 
61.3) 
61.4) 
61.5) 
61.6) 
61.7) 
61.8) 
61.9) 
61.10) 
61.11) 
61.12) 
62.1) 
62.2) 
62.3) 
62.4) 
62.5) 
62.6) 
62.7) 
62.8) 
62.9) 
63.1) 
63.2) 
63.3) 
63.4) 
63.5) 
63.6) 

63.7) 
63.8) 
63.9) 
63.10) 
63.11) 
64.1) 
64.2) 
64.3) 
64.4) 
64.5) 
64.6) 
64.7) 
65.1) 
65.2) 
65.3) 
65.4)
65.5) 
65.6) 
65.7) 
66.1) 
66.2)
66.3)
66.4) 
66.5)
66.6) 
66.7)
67.1)
67.2)
67.3) 
67.4)
67.5)
67.6) 
67.7)
68.1) 
68.2) 
68.3) 
68.4) 
68.5) 
68.6) 
68.7) 
68.8)
68.9) 
69.1) 

69.2)

69.3) 

69.4) 

69.5) 

69.6) 

70)

1.3. **Quivers with relations of tilted algebras of type E_8.**

1.1) 

1.2) 

1.3)

2.1) 

2.2) 

2.3) 

3.1)

3.2) 

3.3) 

4.1) 

4.2) 

4.3) 

5.1) 

5.2) 

5.3) 

6.1) 

6.2) 

6.3) 

6.4)

6.5) 

6.6) 

6.7) 

7.1) 

7.2) 

7.3) 

7.4) 

7.5) 


7.6)
7.7)
8.1)
8.2)
8.3)
8.4)
8.5)
8.6)
8.7)
9.1)
9.2)
9.3)
9.4)
9.5)
9.6)
9.7)
10.1)
10.2)
10.3)
10.4)
10.5)
10.6)
10.7)
11.1)
11.2)
11.3)
11.4)
12.1)
12.2)
12.3)
12.4)
12.5)
12.6)

12.7) 

12.8)

12.9)

12.10) 

12.11) 

12.12) 

12.13) 

12.14) 

13.1) 

13.2)

13.3)

13.4)

14.1)

15.1)

15.2)

16.1)

16.2) 

16.3) 

16.4) 

17.1)

17.2) 

17.3) 

17.4)

18.1)

18.2) 

18.3) 

18.4) 

18.5) 

19.1)

19.2)

19.3) 

19.4) 

19.5) 

20.1) 

20.2)

20.3) 


20.4)
20.5)
20.6)
21.1)
21.2)
21.3)
21.4)
21.5)
21.6)
21.7)
22.1)
22.2)
22.3)
23.1)
23.2)
23.3)
24.1)
24.2)
24.3)
25.1)
25.2)
25.3)
26.1)
26.2)
26.3)
26.4)
26.5)
26.6)
26.7)
27.1)
27.2)
27.3)
27.4)
27.5)

27.6) 

27.7) 

28.1) 

28.2) 

28.3) 

28.4)

28.5) 

28.6) 

29.1) 

29.2) 

29.3) 

29.4) 

29.5) 

29.6)

30.1) 

30.2) 

30.3) 

30.4) 

30.5) 

30.6) 

30.7) 

31.1) 

31.2) 

31.3) 

31.4)

31.5)

31.6) 

32.1) 

32.2) 

32.3) 

32.4) 

32.5) 

32.6) 

32.7) 

33.1) 

33.2) 

33.3) 

33.4)

33.5)

33.6) 

33.7)

34.1) 

34.2) 

34.3) 

34.4) 

34.5)

34.6) 

35.1)

35.2) 

35.3) 

35.4) 

35.5) 

36.1) 

36.2) 

36.3) 

36.4) 

36.5) 

37.1) 

37.2) 

37.3)

37.4)

37.5)

38.1) 

38.2) 

38.3) 

38.4) 

38.5) 

39.1) 

39.2) 

39.3) 

39.4) 

39.5) 

40.1) 

40.2) 

40.3) 

40.4) 

40.5) 

41.1) 
41.2) 
41.3) 
41.4) 
41.5) 
42.1) 
42.2) 
42.3)
42.4) 
42.5) 
43.1) 
43.2) 
43.3) 
43.4) 
43.5) 
44.1) 
44.2) 
44.3) 
44.4) 
44.5) 
45.1) 
45.2) 
45.3) 
45.4) 
45.5) 
46.1) 
46.2) 
46.3) 
46.4) 
46.5) 
46.6) 
47.1) 
47.2) 
47.3) 
47.4) 
47.5) 
47.6) 
48.1) 
48.2) 
48.3) 
48.4) 
48.5) 

48.6) 

48.7) 

48.8) 

49.1) 

49.2) 

49.3) 

49.4) 

49.5) 

49.6) 

49.7) 

49.8) 

50.1) 

50.2) 

50.3) 

50.4) 

50.5) 

50.6) 

51.1) 

51.2) 

51.3) 

51.4) 

51.5) 

51.6) 

52.1) 

52.2) 

52.3) 

52.4) 

53.1) 

53.2) 

53.3) 

53.4) 

53.5) 

53.6) 

53.7) 

53.8) 

54.1) 

54.2) 

54.3)

54.4) 

54.5) 

54.6) 

54.7) 

54.8) 

54.9) 

54.10) 

55.1) 

55.2) 

55.3) 

55.4) 

55.5) 

55.6) 

55.7) 

55.8) 

55.9) 

55.10) 

56.1) 

56.2) 

56.3) 

56.4) 

56.5) 

56.6) 

56.7) 

56.8) 

56.9) 

56.10) 

57.1) 

57.2) 

57.3)

57.4) 

57.5) 

57.6) 

57.7) 

57.8) 

57.9) 

57.10) 

58.1) 

58.2) 

58.3) 

58.4) 

58.5) 

58.6) 

58.7) 

58.8) 

58.9) 

59.1) 

59.2) 

59.3)

59.4) 

59.5) 

59.6) 

59.7) 

59.8) 

59.9) 

60.1) 

60.2) 

60.3) 

60.4) 

60.5) 

60.6) 

60.7) 

60.8) 

60.9) 

60.10) 

61.1) 

61.2) 

61.3) 

61.4) 

61.5) 

61.6) 

61.7) 

61.8) 

61.9) 

62.1) 

62.2) 

62.3) 

62.4) 
62.5)
62.6) 
62.7) 
62.8) 
62.9) 
62.10) 
63.1) 
63.2) 
63.3) 
63.4)
63.5) 
63.6) 
63.7) 
63.8) 
63.9) 
63.10) 
64.1) 
64.2) 
64.3) 
64.4) 
64.5) 
64.6) 
64.7) 
64.8) 
64.9) 
65.1) 
65.2) 
65.3) 
65.4) 
65.5) 
65.6) 
65.7) 
65.8) 
65.9) 
66.1) 
66.2) 
66.3) 
66.4) 
66.5) 
66.6) 
66.7) 
66.8) 
66.9) 
67.1) 
67.2) 

67.3) 
67.4) 
67.5) 
67.6) 
67.7) 

67.8)
67.9) 
68.1) 
68.2) 

68.3) 
68.4) 
68.5) 
68.6) 
68.7) 

68.8) 
68.9) 
69.1) 
69.2) 

69.3) 
69.4) 
69.5) 

69.6) 
69.7) 
69.8) 
69.9) 

69.10) 
69.11) 
69.12) 

70.1) 
70.2) 
70.3) 

70.4) 
70.5) 
70.6) 

70.7) 
70.8) 
70.9) 
70.10) 

71.1) 

71.2) 

71.3) 

71.4) 

71.5) 

71.6) 

71.7) 

71.8) 

71.9) 

71.10) 

71.11) 

71.12) 

71.13) 

71.14) 

72.1) 

72.2) 

72.3) 

72.4) 

72.5) 

72.6) 

72.7) 

72.8) 

72.9) 

72.10) 

72.11) 

72.12) 

72.13) 

72.14) 

72.15) 

73.1) 

73.2) 

73.3) 

73.4) 

73.5) 

73.6) 

73.7) 

73.8) 

73.9) 

73.10) 

74.1) 

74.2) 

74.3) 

74.4) 
74.5) 
74.6) 
74.7) 

74.8) 
74.9) 
74.10) 
75.1) 
75.2) 

75.3) 
75.4) 
75.5) 
75.6) 

75.7) 
75.8) 
75.9) 
75.10) 

76.1) 
76.2) 
76.3) 
76.4) 

76.5) 
76.6) 
76.7 
76.8) 
76.9) 

76.11) 
77.1) 
77.2) 
77.3) 
77.4) 

77.5) 
77.6) 
77.7) 
77.8) 

77.9) 
77.10) 
77.11) 
78.1) 

78.2) 
78.3) 
78.4) 
78.5) 
78.6) 

78.7) 

78.8) 

78.9) 

78.10) 

78.11) 

79.1) 

79.2) 

79.3) 

79.4) 

79.5) 

79.6) 

79.7) 

79.8) 

79.9) 

79.10) 

79.11) 

79.12) 

80.1) 

80.2) 

80.3) 

80.4) 

80.5) 

80.6) 

80.7) 

80.8) 

80.9) 

80.10) 

80.11) 

81.1) 

81.2) 

81.3) 

8.4) 

81.5) 

81.6) 

81.7) 

81.8) 

82.1) 

82.2) 

82.3) 

82.4) 

82.5) 

82.6) 

83.1) 

83.2) 

83.3) 

83.4) 

83.5) 

83.6) 

83.7) 

83.8) 

83.9) 

83.10) 

84.1) 

84.2)

84.3)

84.4) 

84.5) 

84.6)

84.7) 

84.8) 

85.1) 

85.2)

85.3) 

85.4) 

85.5) 

85.6) 

85.7) 

85.8) 

86.1) 

86.2) 

86.3) 

86.4) 

86.5) 

87.1) 

87.2) 

87.3) 

87.4)

88.1)

88.2) 

88.3) 

88.4) 

88.5) 

88.6) 

88.7) 

88.8)

89.1)

89.2)

89.3) 

89.4) 

89.5)

89.6) 

89.7) 

89.8)

90.1)

90.2) 

90.3) 

90.4) 

90.5)

90.6) 

90.7) 

90.8) 

91.1)

91.2)

91.3) 

91.4)

91.5) 

91.6) 

91.7)

91.8) 

92.1) 

92.2)

92.3) 

92.4)

92.5) 

93.1) 

93.2)

93.3)

93.4) 

93.5) 

94.1) 

94.2) 

94.3)

94.4) 

94.5)

95.1) 

95.2) 

95.3)

95.4) 

95.5) 

96.1) 

96.2) 

96.3) 

96.4)

96.5) 

97.1) 

97.2) 

97.3) 

97.4) 

97.5) 

98.1) 

98.2) 

98.3) 

98.4) 

98.5)

99.1) 

99.2) 

99.3) 

99.4)

99.5) 

100.1) 

100.2) 

100.3) 

100.4) 

100.5) 

100.6) 

100.7) 

101.1) 

101.2) 

101.3) 

101.4) 

101.5) 

101.6) 

101.7) 

102.1) 

102.2) 

102.3) 

102.4) 

102.5) 

102.6) 

102.7)

103.1) 

103.2) 

103.3) 

103.4)

103.5) 

103.6) 

103.7) 

104.1) 

104.2) 

104.3) 

104.4) 

104.5) 

104.6) 

104.7) 

105.1) 

105.2) 

105.3)

105.4) 

105.5) 

105.6) 

105.7)

106.1) 

106.2) 

106.3) 

106.4) 

106.5)

106.6) 

106.7) 

107.1) 

107.2) 

107.3) 

107.4) 

107.5) 

107.6) 

108.1) 

108.2) 

108.3) 

108.4) 

108.5) 

108.6) 

109.1) 

109.2) 

109.3) 

109.4) 

109.5) 

109.6) 

110.1) 

110.2) 

110.3) 

111.1) 

111.2) 

111.3) 

111.4) 

111.5) 

111.6) 

111.7) 

111.8) 


112.1)
112.2)
112.3)
112.4)
112.5)
112.6)
112.7)
112.8)
113.1)
113.2)
113.3)
113.4)
113.5)
113.6)
113.7)
113.8)
114.1)
114.2)
114.3)
114.4)
114.5)
114.6)
114.7)
114.8)
115.1)
115.2)
115.3)
115.4)
115.5)
115.6)
115.7)
115.8)
116.1)
116.2)
116.3)
116.4)
116.5)
116.6)

116.7)

116.8) 

116.9) 

117.1) 

117.2) 

117.3) 

117.4)

117.5)

117.6) 

117.7) 

117.8)

117.9) 

118.1) 

118.2) 

118.3) 

118.4) 

118.5)

118.6)

118.7) 

118.8)

118.9) 

118.10)

119.1) 

119.2) 

119.3)

119.4)

119.5) 

119.6) 

119.7) 

119.8)

119.9) 

120.1) 

120.2) 

120.3) 

120.4) 

120.5) 

120.6) 

120.7) 
120.8) 
120.9) 
121.1) 
121.2) 
121.3) 
121.4) 
121.5) 
121.6) 
121.7) 
121.8) 
121.9) 
122.1) 
122.2) 
122.3) 
122.4) 
122.5) 
122.6) 
122.7) 
122.8) 
122.9) 
123.1) 
123.2) 
123.3) 
123.4) 
123.5) 
123.6) 
123.7) 
123.8) 
123.9)
124.1) 
124.2) 
124.3) 
124.4) 
124.5) 
124.6) 
124.7) 
124.8) 

124.9) 

125.1) 

125.2) 

125.3) 

125.4) 

125.5) 

125.6) 

125.7) 

126.1) 

126.2) 

126.3) 

126.4) 

126.5) 

126.6)

126.7) 

127.1) 

127.2) 

127.3) 

127.4) 

127.5) 

127.6) 

127.7) 

127.8) 

127.9) 

127.10) 

128.1) 

128.2) 

128.3) 

128.4) 

128.5) 

128.6) 

128.7) 

128.8) 

128.9) 

128.10) 

129.1) 

129.2) 

129.3) 

129.4) 

129.5) 

129.6) 

129.7) 

129.8) 

129.9) 

129.7) 

130.1) 

130.2) 

130.3) 

130.4) 

130.5) 

130.6) 

131.1) 

131.2) 

131.3) 

131.4) 

131.5) 

131.6)

132.1) 

132.2) 

132.3) 

132.4) 

132.5) 

132.6) 

133.1) 

133.2) 

133.3) 

133.4) 

133.5) 

133.6) 

134.1) 

134.2) 

134.3) 

134.4) 

134.5) 
134.6) 
134.7) 
134.8) 
134.9) 
134.10) 
135.1) 
135.2) 
135.3) 
135.4) 
135.5) 
135.6) 
135.7) 
135.8) 
135.9) 
135.10) 
136.1) 
136.2) 
136.3) 
136.4) 
136.5) 
136.6) 
136.7) 
136.8) 
137.1) 
137.2) 
137.3) 
137.4) 
137.5) 
137.6) 
137.7) 
137.8) 
138.1) 
138.2)
138.3) 
138.4) 

138.5) 
138.6) 
138.7) 
138.8) 
138.9) 
138.10) 
139.1) 
139.2) 
139.3) 
139.4) 
139.5) 
139.6) 
139.7) 
139.8) 
140.1) 
140.2) 
140.3) 
140.4) 
140.5) 
140..6) 
140.7) 
140.8) 
140.9)
141.1) 
141.2) 
141.3) 
141.4) 
141.5) 
141.6) 
141.7) 
141.8) 
141.9) 
142.1) 
142.2) 
142.3) 
142.4) 
142.5) 

142.6) 
142.7) 
142.8) 
142.9) 

143.1)
143.2) 
143.3) 
143.4) 

143.5) 
143.6) 
143.7) 
143.8) 

143.9) 
144.1)
144.2) 

144.3) 
144.4)
144.5)

144.6)
144.7)
144.8) 

144.9) 
144.10)
144.11)

145.1) 
145.2)
145.3) 

145.3)
145.4)
145.5)

145.7) 
145.8) 
145.9) 

145.10)
145.11) 
145.12) 

145.13)
146.1) 
146.2) 

146.3) 

146.4) 

146.5) 

146.6) 

146.7) 

146.8) 

146.9) 

146.10)

146.11)

146.12)

147.1) 

147.2)

147.3) 

147.4) 

147.5) 

147.6) 

147.7) 

147.8) 

147.9) 

147.10) 

147.11) 

147.12) 

147.13)

148.1) 

148.2) 

148.3) 

148.4) 

148.5) 

148.6) 

148.7) 

148.8) 

148.9) 

148.10) 

148.11) 

149.1) 

149.2) 

149.3) 

149.4) 

149.5) 

149.6) 

149.7) 

149.8) 

149.9) 

149.10) 

149.11) 

150.1) 

150.2) 

150.3) 

150.4) 

150.5) 

150.6) 

150.7) 

150.8) 

150.9) 

150.10) 

150.11) 

151.1) 

151.2) 

151.3) 

151.4) 

151.5) 

151.6) 

151.7) 

151.8) 

151.9) 

151.10) 

152.1) 

152.2) 

152.3) 

152.4) 

152.5) 

152.6) 

152.7) 

152.8) 

152.9) 

152.10) 

152.11) 

152.12) 

152.13) 

153.1) 

153.2) 

153.3) 

153.4) 

153.5) 

153.6) 

153.7) 

153.8) 

153.9) 

153.10) 

153.11) 

153.12) 

153.13) 

153.14) 

154.1) 

154.2) 

154.3) 

154.4) 

154.5) 

154.6) 

154.7) 

154.8) 

154.9) 

154.10) 

155.1) 

155.2) 

155.3) 

155.4) 

155.5) 

155.6) 

155.7) 

155.8) 

155.9) 

155.10) 

155.11) 

155.12) 

155.13) 

156.1) 

156.2) 

156.3) 

156.4) 

156.5) 

156.6) 

156.7) 

156.8) 

156.9) 

156.10) 

157.1) 

157.2) 

157.3) 

157.4) 

157.5) 

157.6) 

157.7) 

157.8) 

158.1) 

158.2) 

158.3) 

158.4) 

158.5) 

158.6) 

158.7) 
158.8) 
158.9) 
158.10) 
158.11) 
159.1)
159.2) 
159.3) 
159.4) 
159.5) 
159.6) 
159.7) 
159.8) 
159.9) 
159.10)
160.1) 
160.2) 
160.3) 
160.4) 
160.5) 
160.6) 
160.7) 
160.8) 
160.9) 
160.10) 
160.11)
160.12) 
161.1) 
161.2) 
161.3)
161.4)
161.5) 
162.1) 
162.2) 
162.3)
162.4)
162.5) 

163.1) 

163.2) 

163.3) 

163.4) 

163.5) 

163.6) 

163.7) 

163.8)

163.9) 

163.10) 

164.1) 

164.2) 

164.3) 

164.4) 

164.5) 

165.1) 

165.2) 

165.3) 

165.4) 

165.5) 

166.1) 

166.2) 

166.3) 

166.4) 

166.5) 

166.6) 

166.7) 

166.8) 

166.9) 

167.1) 

167.2) 

167.3) 

167.4) 

167.5) 

167.6) 

167.7) 

167.8) 

167.9) 

168.1) 

168.2) 

168.3) 

168.4) 

168.5) 

168.6)

168.7) 

168.8) 

168.9) 

169.1) 

169.2) 

169.3) 

169.4) 

169.5) 

169.6)

169.7) 

169.8) 

169.9) 

170.1) 

170.2) 

170.3) 

170.4) 

170.5) 

170.6) 

170.7) 

170.8) 

170.9)

170.10)

171.1) 

171.2) 

171.3)

171.4)

171.5) 

171.6)

171.7) 

171.8)

171.9)

171.10) 

171.11) 

171.12) 

171.13) 

171.14) 

172.1)

172.2) 

171.3) 

171.4) 

172.5)

172.6)

172.7)

172.8) 

172.9) 

172.10) 

172.11) 

173.1) 

173.2) 

173.3) 

173.4) 

173.5) 

173.6) 

174.1) 

174.2) 

174.3) 

174.4) 

174.5) 

174.6) 

174.7) 

174.8) 

174.9) 

174.10) 

174.11) 

174.12) 

175.1) 

175.2) 

175.3) 

175.4) 

175.5) 

175.6) 

175.7) 

176.1) 

176.2) 

176.3) 

176.4) 

176.5) 

176.6) 

176.7) 

176.8) 

176.9) 

177.1) 

177.2) 

177.3) 

177.4) 

177.5) 

177.6) 

177.7) 

177.8) 

177.9) 

178.1) 

178.2) 

178.3) 

178.4) 

178.5) 

178.6) 

178.7) 

178.8) 

178.9) 

178.10) 

179.1) 

179.2)

179.3) 

179.4) 

179.5)

179.6) 

179.7) 

180.1) 

180.2) 

180.3) 

180.4) 

180.5) 

180.6) 

180.7) 

181.1) 

181.2) 

181.3) 

181.4)

181.5) 

181.6) 

181.7) 

182.1) 

182.2) 

181.3)

182.4) 

182.5) 

182.6) 

182.7) 

183.1)

183.2) 

183.3)

183.4) 

183.2) 

184.1)

184.2) 

184.3)

184.4)

184.5)

185.1)

185.2)

185.3)

185.4) 

185.5) 

186.1) 

186.2) 

186.3) 

186.4) 

186.5) 

186.6) 

186.7) 

186.8) 

187.1) 

187.2) 

187.3) 

187.4) 

187.5) 

187.6) 

187.7) 

187.8) 

188.1) 

188.2) 

188.3) 

188.4) 

188.5) 

188.6) 

188.7) 

188.8) 

188.9) 

188.10) 

188.11) 

189.1) 

189.2) 

189.3) 

189.4) 

189.5) 

188.6) 

189.7) 

190.1) 

190.2) 

190.3) 

190.4) 

190.5) 

190.6) 

190.7) 

190.8) 

190.9) 

190.10) 

190.11) 

191.1) 

191.2) 

191.3) 

191.4) 

191.5) 

191.6) 

191.7) 

191.8) 

191.9) 

192.1) 

192.2) 

192.3) 

192.4) 

192.5) 

192.6) 

192.7) 

192.8) 

192.9) 

193.1) 

193.2) 

193.3) 

193.4) 
193.5) 
193.6) 
193.7) 
193.8) 
193.9) 
194.1) 
194.2) 
194.3) 
194.4) 
194.5) 
194.6) 
194.7) 
194.8) 
194.9) 
194.10) 
194.11) 
194.12) 
195.1) 
195.2) 
195.3) 
195.4) 
195.5) 
195.6) 
195.7) 
195.8) 
195.9) 
195.10) 
196.1) 
196.2) 
196.3) 
196.4) 
196.5) 

196.6) 

196.7) 

196.8) 

196.9) 

196.10) 

197.1) 

197.2)

197.3) 

197.4) 

197.5)

197.6) 

197.7)

197.8) 

197.9) 

198.1) 

198.2)

198.3)

198.4) 

198.5)

198.6) 

198.7) 

198.8) 

198.9) 

198.10)

198.11)

199.1) 

199.2) 

199.3) 

199.4)

199.5) 

199.6) 

199.7) 

199.8) 

199.9)

199.10)

199.11) 

200.1)

200.2) 

200.3) 

200.4) 

200.5) 

201.1)

201.2)

201.3) 

201.4) 

201.5) 

202.1) 

202.2) 

202.3) 

202.4) 

202.5) 

202.6) 

202.7) 

202.8) 

202.9) 

202.10) 

203.1) 

203.2) 

203.3) 

203.4) 

203.5) 

203.6) 

203.7) 

203.8) 

203.9) 

203.10) 

204.1) 

204.2) 

204.3)

204.4) 

204.5)

204.6)

204.7)

204.8) 

204.9) 

204.10) 

205.1) 

205.2) 

205.3) 

205.4) 

205.5) 

205.6) 

205.7) 

205.8) 

205.9) 

206.1) 

206.2) 

206.3) 

206.4) 

206.5) 

206.6) 

206.7) 

206.8) 

206.9) 

207.1) 

207.2) 

207.3) 

207.4) 

207.5) 

207.6) 



2. Cluster tilted algebras of type E_p

2.1. **Quivers with relations of cluster tilted algebras of type E_6.**



11)  12)  13)  14)  15) 

16)

2.2. **Quivers with relations of cluster tilted algebras of type E_7.**

1)  2)  3)  4) 

5)  6)  7)  8) 

9) 10) 11)  12) 

13)  14)  15)  16) 

17)  18)  19)  20) 

21)  22)  23)  24)  25) 

26)  27)  28)  29)  30) 

31)  32) 33)  34)  35) 



2.3. **Quivers with relations of tilted algebras of type E_8.**




4)
5)
6)
7)
8)
9)
10)
11)
12)
13)
14)
15)
16)
17)
18)
19)
20)
21)
22)
23)
24)
25)
26)
27)
28)
29)
30)
31)
32)
33)
34)
35)
36)
37)
38)
39)
40)
41)
42)
43)
44)
45)
46)
47)
48)
49)
50)
51)
52)

53) 

54) 

55) 

56) 

57) 

58) 

59) 

60) 

61) 

62) 

63) 

64) 

65) 

66) 

67) 

68) 

69) 

70) 

71) 

72) 

73) 

74) 

75) 

76) 

77) 

78) 

79) 

80) 

81) 

82) 

83) 

84) 

85) 

86) 

87) 

88) 

89) 

90) 

91) 

92) 

93) 

94) 

95) 

96) 

97) 

98) 

99) 

100) 

101)

102) 
103) 
104) 
105) 
106) 
107) 
108) 
109) 
110) 
111) 
112) 
113) 
114) 
115) 
116) 
117) 
118) 
119) 
120) 
121) 
122) 
123) 
124) 
125) 
126) 
127) 
128) 
129) 
130) 
131) 
132) 
133) 
134) 
135) 
136) 
137) 
138) 
139) 
140) 
141) 
142) 
143) 
144) 

145) 
146) 
147) 
148) 

149) 
150) 
151) 
152) 
153) 

154) 
155) 
156) 
157) 

158) 
159) 
160) 
161) 

162) 
163) 
164) 
165) 

166) 
167) 
168) 
169) 
170) 

171)
172)
173) 
174) 

175) 
176) 
177) 
178) 
179) 

180) 
181) 
182) 
183) 
184) 

185) 
186) 
187) 
188) 
189) 



References

[1] N. Bordino, E. Fernández, S. Trepode. *On the quiver with relations of a quasitilted algebra and applications*. In preparation.

Natalia Bordino, Departamento de Matemática, Facultad de Ciencias Exactas y Naturales, Funes 3350, Universidad Nacional de Mar del Plata, 7600 Mar del Plata, Argentina.

E-mail address: bordino@mdp.edu.ar

Elsa Fernández, Facultad de Ingeniería, Universidad Nacional de la Patagonia San Juan Bosco, 9120 Puerto Madryn, Argentina.

E-mail address: elsafer9@gmail.com

Sonia Trepode, Departamento de Matemática, Facultad de Ciencias Exactas y Naturales, Funes 3350, Universidad Nacional de Mar del Plata, 7600 Mar del Plata, Argentina.

E-mail address: strepode@mdp.edu.ar